

(3.4) Estimate if fusion of deuterium into helium releases more or less energy per gram of material consumed than the fission of uranium.

First we define the Avogadro number, N_A , and the electron volt, eV.

$$N_A := 6.0221367 \cdot 10^{23} \cdot \text{mole}^{-1} \quad \text{eV} := 1.60217733 \cdot 10^{-19} \cdot \text{joule} \quad \text{MeV} := 10^6 \cdot \text{eV}$$

The fission energy of a ^{235}U atom, Q_U , is given by eqn. (19.1):

$$Q_U := 200 \cdot \text{MeV}$$

The atomic weight of ^{235}U , Mw_U , and of ^4He , Mw_{He} , can be approximated as follows:

$$Mw_U := 235 \cdot \text{gm} \cdot \text{mole}^{-1} \quad Mw_{He} := 4 \cdot \text{gm} \cdot \text{mole}^{-1}$$

$$Q_{fiss} := \frac{Q_U \cdot N_A}{Mw_U} \quad Q_{fiss} = 5.125 \cdot 10^{23} \cdot \text{MeV} \cdot \text{gm}^{-1}$$

$$EB_{He} := 28.30 \cdot \text{MeV} \quad EB_D := 2.22 \cdot \text{MeV}$$

$$Q_{He} := EB_{He} - 2 \cdot EB_D \quad Q_{He} = 23.86 \cdot \text{MeV}$$

Neglect the very small mass difference between fuel as 2 D and helium in the calculation of the energy content:

$$Q_{fus} := \frac{Q_{He} \cdot N_A}{Mw_{He}} \quad Q_{fus} = 3.592 \cdot 10^{24} \cdot \text{MeV} \cdot \text{gm}^{-1}$$

$$\frac{Q_{fus}}{Q_{fiss}} = 7.009$$

Thus fusion gives about 7 times more energy per g of fuel