(13.8) A thin Au target is irradiated with a beam of 800 MeV  $^{16}\text{O}^{2}$ + ions during 2 hours. After the irradiaten a faraday cup (with current integrator) behind the target showed a total accumulated charge of 92.3  $\mu$ C. What was the average beam intensity (oxygen ions per second)? Consider charge-stripping in the target.

Definitions of units and constants:

$$q_e := 1.6021773 \cdot 10^{-19} \cdot coul$$
  $N_A := 6.022137 \cdot 10^{23} \cdot mole^{-1}$   $c_{light} := 299792458 \cdot m \cdot sec^{-1}$   $m_e := 9.109390 \cdot 10^{-31} \cdot kg$   $M_{16O} := 15.994915 \cdot gm \cdot mole^{-1}$   $MeV := 1.6021773 \cdot 10^{-13} \cdot joule$   $m_{OOion} := \frac{M_{16O}}{N_A} - 2 \cdot m_e$  The rest mass of  $^{16O2+}$  ions

Then compute the mass of <sup>16</sup>O<sup>2+</sup> ions at 800 MeV kinetic energy by using eqn. (4.21)

$$E_{Oion} = 800 \cdot MeV$$
  $m_{Oion} = m_{OOion} + \frac{E_{Oion}}{c_{light}^2}$ 

Then use eqn (4.19) to compute  $\beta$  with  $\beta = (v/c)^2$  and  $\nu$  from  $\nu = \beta *c$ :

$$\beta := \sqrt{1 - \left(\frac{m_{OOion}}{m_{Oion}}\right)^2} \qquad \beta = 0.315 \qquad v_{Oion} := \beta \cdot c_{light}$$

When the velocity is known, we can use eqn. (13.2) to compute the effective charge after passage of the target foil:

$$Z := 8$$
  $k := 3.6 \cdot 10^6 \cdot m \cdot sec^{-1}$   $Q := 92.3 \cdot 10^{-6} \cdot coul$ 

$$z_{eff} := Z \cdot \left[ 1 + \left( \frac{v \cdot Oion}{k \cdot Z^{0.45}} \right)^{-1.67} \right]^{-0.6}$$

$$z_{eff} = 7.904$$

Once we know the effective average charge of an ion hitting the faraday cup, we can compute the number of ions per unit time as follows:

$$t_{irr} = 2 \cdot hr$$
  $I_{ion} = \frac{Q}{t_{irr} z_{eff} q_{e}}$   $I_{ion} = 1.01 \cdot 10^{10} \cdot sec^{-1}$